

Thyroid Synergy™



– Nutritional Support for the Thyroid Gland

By David Brady, ND, DC, CCN, DACBN & Suzanne Copp, MS

THIS INFORMATION IS PROVIDED FOR THE USE OF PHYSICIANS AND OTHER LICENSED HEALTH CARE PRACTITIONERS ONLY. THIS INFORMATION IS INTENDED FOR PHYSICIANS AND OTHER LICENSED HEALTH CARE PROVIDERS TO USE AS A BASIS FOR DETERMINING WHETHER OR NOT TO RECOMMEND THESE PRODUCTS TO THEIR PATIENTS. THIS MEDICAL AND SCIENTIFIC INFORMATION IS NOT FOR USE BY CONSUMERS. THE DIETARY SUPPLEMENT PRODUCTS OFFERED BY DESIGNS FOR HEALTH ARE NOT INTENDED FOR USE BY CONSUMERS AS A MEANS TO CURE, TREAT, PREVENT, DIAGNOSE, OR MITIGATE ANY DISEASE OR OTHER MEDICAL CONDITION.

Like all of our glands, the thyroid needs supportive nutrients from food or supplements to act as raw ingredient precursors for its proper function. Thyroid Stimulating Hormone needs to give the cells a message to synthesize thyroxine. Thyroxine, known as T₄, needs to be converted into active T₃. Cortisol and insulin levels need to be controlled in order for this to happen successfully and precursors and coenzymes must be available.

SUPPORTIVE HERBS: AMERICAN GINSENG ROOT AND COLEUS FORSKOHLII (FORSLEAN®)

The non-stimulating adaptogenic botanical American Ginseng (*Panax quinquefolius*) helps control variations in cortisol. These variations can have profound deleterious effects on peripheral thyroid hormone conversion, as well as serum blood glucose and insulin levels, inducing dysglycemia and laying the foundation for metabolic syndrome.

WHY IS COLEUS FORSKOHLII EXTRACT NECESSARY FOR THYROID SUPPORT?

Coleus forskohlii is necessary for several reasons. Its active constituent forskohlin is a diterpene from the roots of Coleus forskohlii, known to directly activate the adenylate cyclase enzyme and raise cyclic AMP levels in many tissues. This is important because Thyroid Stimulating Hormone (TSH) stimulates all steps in the synthesis and secretion of T₃ and T₄. The first step is the binding of TSH hormone to TSH receptors located on the basal membrane of thyroid epithelial cells, followed by elevation of cyclic AMP levels, and a subsequent cascade of phosphorylation reactions via activation of protein kinase A. The final step is the secretion of thyroid hormones T₄ and T₃ which again could not happen without cAMP.¹⁰ Hypothyroid patients need support in the area of fat metabolism. Coleus is known to reduce adipose assimilation and increase the lipolysis of fats. Coleus aids in the following functions:

- 1) Inhibiting the synthesis of adipose tissue
- 2) Increasing response to lipolytic hormones, which are produced in less quantity as we age
- 3) Utilizing stored fat.¹⁰ It may also play a role in aiding the production of digestive enzymes and supporting the immune system, both of which are helpful for hypothyroid patients.

Supplement Facts

Serving Size 2 capsules
Servings Per Container 60

Amount Per Serving	% Daily Value	Amount Per Serving	% Daily Value		
Vitamin A (as Palmitate)	2000 IU	40%	Chromium	50 mcg	40%
Riboflavin (Vitamin B-2)	5 mg	290%	(TRAACS® Chromium Nicotinate Glycinate Chelate)		
(as Riboflavin-5-Phosphate)			N-Acetyl L-Tyrosine	200 mg	*
Iodine	100 mcg	70%	American Ginseng	200 mg	*
(as Potassium Iodide)			(<i>Panax quinquefolius</i>)(root)		
Zinc	25 mg	170%	[standardized to contain 5% ginsenosides]		
(TRAACS® Zinc Glycinate Chelate)			ForsLean® Extract	50 mg	*
Selenium	50 mcg	70%	(<i>Coleus forskohlii</i>)(root)		
(as Selenomethionine)			[standardized to contain 20% forskohlin]		
Copper	500 mcg	25%			
(TRAACS® Copper Glycinate Chelate)					
Manganese	5 mg	250%			
(TRAACS® Manganese Glycinate Chelate)					

*Daily Value not established.

Other Ingredients: Microcrystalline cellulose, vegetable stearate, rice flour.

TYROSINE, COPPER AND COLEUS FORSKOHLII

Cyclic AMP, copper and tyrosine are involved in the production of "feel good" catecholamines such as dopamine and norepinephrine. This could be very helpful since hypothyroid patients are so susceptible to depression.¹⁰ Cyclic AMP is also a key player in regulating steroid hormone synthesis. These hormones include estrogen, progesterone and testosterone and depend on cholesterol for their making.¹⁰ Women with hypothyroidism can have very severe PMS.

CHROMIUM, ZINC AND MANGANESE

These minerals are needed for thyroid support to help control insulin since insulin blocks phosphorylation and therefore opposes the action of epinephrine and can impede thyroid hormone production.¹⁰ Chromium also helps to control cholesterol levels that often elevate in hypothyroid patients as well as control blood sugar levels which are challenged by poor adrenal and thyroid gland health.

N-ACETYL L-TYROSINE AND IODINE

Tyrosine and iodine are intimately involved in the synthesis of Thyroxine. N-acetyl L-tyrosine is much more stable and better absorbed than L-tyrosine. Vitamin B2 supports these enzymatic conversions. A lack of iodine is a known contributing cause of goiter. Potassium iodide has been shown to protect the thyroid gland from radiation damage.¹ Potassium levels are often low in patients with poor thyroid and poor adrenal function. Potassium is necessary for getting thyroid hormone into our cells and is very energizing.

SELENIUM

Selenium is a key cofactor in the enzymatic conversion (ID-I enzyme) of T₄, thyroxine hormone, to the more active T₃, triiodothyronine. Selenium also protects the thyroid gland from mercury damage and from free radical damage caused by chemicals and heavy metals. Selenium helps to make the powerfully protective enzyme glutathione peroxidase.

One study looked at how selenium deficiency affected ID-I enzyme (I iodothyronine 5'-deiodinase) levels in the body. Selenium deficient rats had the following results: 1.) Hepatic and renal ID-I activities decreased by 60% and 50% respectively, serum levels of T₃ reduced 36%, and T₄ increased by 32%. 2.) GSH-Px (glutathione peroxidase) activities in the whole blood and liver decreased by 61% and 82% respectively. LPO (lipid peroxide) concentrations in serum and liver increased by 53% and 40% respectively.³ The results of a Graves' disease study clearly indicated that supplementation with antioxidants in the treatment of Graves' disease is justified, particularly those containing selenium.⁵

VITAMIN A, ZINC AND SELENIUM

Studies tell us that TTR (Transthyretin), an extracellular transfer protein in humans, is responsible for binding both T₄ and T₃ with a preference for T₄. This protein also binds vitamin A, known to be important for thyroid function. TTR is an extracellular transport protein involved in the distribution of thyroid hormones and vitamin A. Hypothyroid patients often have a yellow-orange coloring to their skin because of very poor conversion of beta-carotene into vitamin A.⁶ Zinc is needed for this conversion. Significantly lower levels of zinc and selenium are found in the thyroids of patients with thyroid cancer, as compared to other thyroid patients.⁷ Zinc is likely as potent a thyroid antioxidant as selenium. Both of these minerals protect against heavy metals.

Q: CAN SELENIUM SUPPLEMENTATION HELP THYROIDITIS?

A: Thyroiditis is inflammation of the thyroid gland. The most common type is autoimmune thyroiditis, in which, for complicated reasons, the body attacks the thyroid gland, decreasing its production of thyroid hormones.

Selenium appears to have two essential functions related to thyroid health. First, it is a component of the detoxifying enzyme glutathione peroxidase. If selenium is unavailable, this enzyme does not work as effectively. Since glutathione peroxidase helps to quench free radicals, lowered activity may result in increased free-radical generation and increased damage to an already inflamed thyroid gland. Second, selenium is a key cofactor in the enzymatic conversion of thyroid hormone thyroxine (T₄) to the more active triiodothyronine (T₃), making selenium important for healthy thyroid function. In two recent studies, researchers tested this idea by giving subjects T₄ with or without selenium supplements. In both studies the combined supplementation of T₄ and selenium, compared with T₄ only, was beneficial in decreasing the autoimmune-induced inflammation as measured by antibodies against thyroid peroxidase, particularly in those with very high antibody levels.¹² These two studies do show selenium is helpful and well-tolerated at the dose of 200 mcg daily.^{11,12}

References

1. Cancer. 2003 Jul 15;98(2):389-96. Improved radiation protection of the thyroid gland with thyroxine, methimazole, and potassium iodide during diagnostic and therapeutic use of radiolabeled metaiodobenzylguanidine in children with neuroblastoma. van Santen HM, de Kraker J, van Eck BL, de Vijlder JJ, Vulmsa T.
2. Ter Arkh. 2003;75(10):54-7. Iodine prophylaxis in pregnant women living in conditions of goiter endemia.
3. Chin Med J (Engl). 1998 Sep;111(9):854-7. Changes of peripheral tissue thyroid hormone metabolism in rats fed with selenium- and vitamin E-deficient artificial semisynthetic diet. Yue L, Wang F, Li G.
4. J Biol Chem. 2004 Apr 13. High-resolution crystal structures of piscine transthyretin reveal different binding modes for triiodothyronine and thyroxine. Eneqvist T, Lundberg E, Karlsson A, Huang S, Santos CR, Power DM, Sauer-Eriksson EA.
5. Clin Chim Acta. 2004 Mar;341(1-2):55-63. Supplementation with antioxidants in the treatment of Graves' disease; the effect on glutathione peroxidase activity and concentration of selenium. Vraa VB, Skreb F, Cepelak I, Romic Z, Mayer L.
6. Acta Med Austriaca. 1993;20(1-2):17-20. Beta-carotene, vitamin A and carrier proteins in thyroid diseases. Aktuna D, Buchinger W, Langsteiger W, Meister E, Sternad H, Lorenz O, Eber O.
7. Biol Trace Elem Res. 2003 Summer;93(1-3):9-18. Copper, zinc, and selenium in whole blood and thyroid tissue of people with various thyroid diseases. Kucharzewski M, Braziewicz J, Majewska U, Gozdz S.
8. Inorg Chem. 1997 Dec 3;36(25):5777-5784. Weak Interactions in Ternary Copper(II) Complexes with Iodotyrosinates. Biological Significance of the Iodines in Thyroid Hormones. Zhang F, Yajima T, Masuda H, Odani A, Yamauchi O.
9. Biol Trace Elem Res. 1992 Jan-Mar;32:173-85. Clinical implications of trace elements in endocrinology. Neve J.
10. Fifth Edition Textbook of Biochemistry by Thomas Devlin. p. 880, 883
11. Gartner R, et al. Selenium supplementation in patients with autoimmune thyroiditis decreases thyroid peroxidase antibodies concentrations. J Clin Endocrinol Metab 2002 Apr;87(4):1687-91.
12. Duntas LH, et al. Effects of a six-month treatment with selenomethionine in patients with autoimmune thyroiditis. Eur J Endocrinol 2003 Apr;148(4):389-93.